

SCHUR'S OLD DETERMINANT PROVES A BRAND-NEW THEOREM OF GARRETT-ISMAIL-STANTON

HELMUT PRODINGER

ABSTRACT. Garrett, Ismail, and Stanton gave a general formula that contains the Rogers–Ramanujan identities as special cases. We show how easy this is when using a determinant that Schur introduced in 1917.

In a recent difficult paper [2] Garrett, Ismail, and Stanton prove, amongst many other things, the following generalization of the celebrated Rogers–Ramanujan identities:

$$\sum_{n \geq 0} \frac{q^{n^2+mn}}{(1-q)(1-q^2) \dots (1-q^n)} = (-1)^m q^{-\binom{m}{2}} E_{m-2} \prod_{n \geq 0} \frac{1}{(1-q^{5n+1})(1-q^{5n+4})} \\ - (-1)^m q^{-\binom{m}{2}} D_{m-2} \prod_{n \geq 0} \frac{1}{(1-q^{5n+2})(1-q^{5n+3})}, \quad (1)$$

with the *Schur* polynomials, defined by

$$\begin{aligned} D_m &= D_{m-1} + q^m D_{m-2}, & D_0 &= 1, \quad D_1 = 1 + q, \\ E_m &= E_{m-1} + q^m E_{m-2}, & E_0 &= 1, \quad E_1 = 1. \end{aligned}$$

(Another proof, based on generalized Engel expansions, can be found in [1].)

Schur [3] has computed the limits

$$D_\infty = \prod_{n \geq 0} \frac{1}{(1-q^{5n+1})(1-q^{5n+4})}, \quad E_\infty = \prod_{n \geq 0} \frac{1}{(1-q^{5n+2})(1-q^{5n+3})}.$$

The aim of this note is to give a simple proof of this result, based on the following determinant of Schur:

$$\text{Schur}(x) := \begin{vmatrix} 1 & xq^{1+m} & & & & \\ -1 & 1 & xq^{2+m} & & & \\ & -1 & 1 & xq^{3+m} & & \\ & & -1 & 1 & xq^{4+m} & \\ & & & \ddots & \ddots & \ddots \end{vmatrix}.$$

Expanding the determinant with respect to the first column (“top-recursion”) we get

$$\text{Schur}(x) = \text{Schur}(xq) + xq^{1+m} \text{Schur}(xq^2).$$

Date: April 2, 2000.

1991 Mathematics Subject Classification. 05A30.

Key words and phrases. Rogers–Ramanujan identities, Schur’s determinant.

Setting

$$\text{Schur}(x) = \sum_{n \geq 0} a_n x^n,$$

we get, upon comparing coefficients,

$$a_n = q^n a_n + q^{1+m} q^{2n-2} a_{n-1},$$

or

$$a_n = \frac{q^{2n-1+m}}{1 - q^n} a_{n-1}.$$

Since $a_0 = 1$, iteration leads to

$$a_n = \frac{q^{n^2+mn}}{(1-q)(1-q^2) \dots (1-q^n)},$$

and thus the left hand side of (1) can be expressed by $\text{Schur}(1)$.

On the other hand, $\text{Schur}(1)$ is the limit of the *finite* determinants

$$\text{Schur}_n := \begin{vmatrix} 1 & q^{1+m} & & & & \dots \\ -1 & 1 & q^{1+m} & & & \dots \\ & -1 & 1 & q^{2+m} & & \dots \\ & & -1 & 1 & q^{3+m} & \dots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots \\ & & & -1 & 1 & q^{n+m} \\ & & & & -1 & 1 \end{vmatrix}.$$

Expanding this determinant with respect to the last row (“bottom–recursion”) we get

$$\text{Schur}_n = \text{Schur}_{n-1} + q^{n+m} \text{Schur}_{n-2}.$$

We see that the sequences $\langle D_{n+m} \rangle_n$ and $\langle E_{n+m} \rangle_n$ satisfy this recursion, and thus any linear combination; set

$$\text{Schur}_n = \lambda D_{n+m} + \mu E_{n+m}.$$

We can determine the parameters λ and μ using the initial conditions $\text{Schur}_0 = 1$, $\text{Schur}_1 = 1 + q^{1+m}$, which leads to

$$\lambda = \frac{E_m - E_{m-1}}{D_{m-1}E_m - D_mE_{m-1}} = \frac{q^m E_{m-2}}{D_{m-1}E_m - D_mE_{m-1}}$$

and

$$\mu = \frac{D_m - D_{m-1}}{D_mE_{m-1} - D_{m-1}E_m} = \frac{q^m D_{m-2}}{D_mE_{m-1} - D_{m-1}E_m}.$$

We can find the nicer form

$$\lambda = \frac{q^m E_{m-2}}{D_{m-1}E_m - D_mE_{m-1}} = (-1)^m q^{-\binom{m}{2}} E_{m-2},$$

which amounts to prove that

$$D_{m-1}E_m - D_mE_{m-1} = (-1)^m q^{\binom{m+1}{2}}.$$

This however is an easy induction on m , the beginning $m = 0$ being trivial. The induction step goes like this:

$$\begin{aligned} D_m E_{m+1} - D_{m+1} E_m &= D_m (E_m + q^{m+1} E_{m-1}) - (D_m + q^{m+1} D_{m-1}) E_m \\ &= q^{m+1} (D_m E_{m-1} - D_{m-1} E_m) \\ &= -q^{m+1} (-1)^m q^{\binom{m+1}{2}} = (-1)^{m+1} q^{\binom{m+2}{2}}. \end{aligned}$$

Analogously we obtain

$$\mu = \frac{q^m D_{m-2}}{D_m E_{m-1} - D_{m-1} E_m} = (-1)^{m+1} q^{-\binom{m}{2}} D_{m-2}.$$

Hence

$$\text{Schur}_n = (-1)^m q^{-\binom{m}{2}} E_{m-2} D_{n+m} - (-1)^m q^{-\binom{m}{2}} D_{m-2} E_{n+m}.$$

Performing the limit $n \rightarrow \infty$ this turns into

$$\text{Schur}(1) = (-1)^m q^{-\binom{m}{2}} E_{m-2} D_\infty - (-1)^m q^{-\binom{m}{2}} D_{m-2} E_\infty,$$

which is (1).

REFERENCES

- [1] G. Andrews, A. Knopfmacher, and P. Paule. An Infinite Family of Engel Expansions of Rogers-Ramanujan Type. *Adv. in Appl. Math.*, to appear 1999.
- [2] K. Garrett, M. Ismail, and D. Stanton, Variants of the Rogers-Ramanujan identities, *Adv. in Appl. Math.* **23** (1999), 274–299.
- [3] I. Schur. Ein Beitrag zur additiven Zahlentheorie und zur Theorie der Kettenbrüche. *S.-B. Preuss. Akad. Wiss. Phys.-Math. Kl.*, 1917, 302–321, reprinted in I. Schur, *Gesammelte Abhandlungen*, vol. 2, pp. 117–136, Springer, 1973.

HELMUT PRODINGER, THE JOHN KNOPFMACHER CENTRE FOR APPLICABLE ANALYSIS AND NUMBER THEORY, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, P. O. WITS, 2050 JOHANNESBURG, SOUTH AFRICA, EMAIL: helmut@gauss.cam.wits.ac.za,
 HOMEPAGE: <http://www.wits.ac.za/helmut/index.htm>